

## Fixed Points for Multivalued Mappings Defined on Unbounded Sets in Banach Spaces

GIUSEPPE MARINO

*Dipartimento Matematica, Università della Calabria,  
Cosenza, Italy*

*Submitted by Ky Fan*

Received July 24, 1989

We give some fixed points theorems for multivalued nonexpansive and pseudocontractive mappings defined on closed convex unbounded subsets of a Banach space. Some of such results extend to the multivoque case a few of the theorems already known for the univoque case, while some other results improve a few of the theorems already known for the multivoque case. © 1991 Academic Press, Inc.

In this paper we study the existence of fixed points for multivalued mappings  $T: K \rightarrow X$ , where  $K$  is an unbounded closed convex subset of a real Banach space  $X$  and  $T$  is either a nonexpansive or a pseudocontractive mapping (see below for their definitions).

Our investigation is prompted by a work of Kirk–Ray [1] in which similar problems are treated for single-valued mappings in uniformly convex Banach spaces and by two recent papers of Carbone–Marino [2] and Marino–Pietramala [3] in which (always for single-valued case) the structure of some geometric sets in Banach spaces is examined.

The theorem of Lim [4] always assures the existence of a fixed point for nonexpansive mappings  $T: K \rightarrow \mathcal{K}(K)$  from the closed bounded convex subset  $K$  of an uniformly convex Banach space  $X$  into the family of non-empty compact subsets of  $K$ , while it is known that for a wide class of unbounded closed convex sets  $K$  (e.g., subsets of Hilbert spaces which contain an infinite ray) there exist nonexpansive mappings  $T: K \rightarrow \mathcal{K}(K)$  which fail to have a fixed point [1].

Thus it is interesting to investigate the existence of fixed points for non-expansive mappings defined on closed convex unbounded subsets in the “rich” structure of Hilbert spaces too.

A line of research in this direction (for the single-valued case) has been initiated by Goebel–Kuczumov [5] and is expanded in [1, 6, 7, 2, 3].

In these papers it is shown that it is sufficient to assume that  $T$  moves some points in a bounded "direction" for the existence of fixed points.

More precisely in [5] it is proved that if  $K$  is a closed convex subset of  $l_2$  and  $T: K \rightarrow K$  is nonexpansive (single-valued) for which there exists a point  $x \in K$  such that the set

$$LS(x, Tx; K) := \{z \in K : \langle z - x, Tx - x \rangle \geq 0\}$$

is bounded, then  $T$  has a fixed point in  $K$ .

On the same line in [1] it is shown that if  $K$  is an unbounded closed convex subset of an uniformly convex space  $X$  and  $T: K \rightarrow K$  is lipschitzian pseudocontractive (single-valued) mapping for which the set

$$G(x, Tx; K) := \{z \in K : \|z - Tx\| \leq \|z - x\|\}$$

is bounded for some  $x \in K$ , then  $T$  has a fixed point in  $K$ .

The present paper seems to be the first to extend to Banach spaces and multivalued mappings the ideas contained (for single-valued case) in [5, 1].

## 1. NOTATION AND DEFINITIONS

In this section we introduce some necessary notation and definitions.

Let  $X$  be a real Banach space and let  $K$  be a nonempty convex subset of  $X$ . We denote by  $\mathcal{K}(X)$  (resp.  $\mathcal{K}(K)$ ) the family of nonempty compact subsets of  $X$  (resp.  $K$ ) and by  $\mathcal{CB}(X)$  the family of nonempty bounded closed subsets of  $X$ .

For any  $A \in \mathcal{CB}(X)$  we note with  $\overline{\text{co}}(A)$  the closed convex hull of  $A$  and for any  $A, B \in \mathcal{CB}(X)$  the Hausdorff metric  $H$ , induced by the norm of  $X$ , is defined as

$$H(A, B) := \max(\sup_{b \in B} d(b, A), \sup_{a \in A} d(a, B)),$$

where  $d(x, Y) := \inf\{\|x - y\| : y \in Y\}$ .

A multivalued mapping  $T: K \rightarrow \mathcal{CB}(X)$  is said to be lipschitzian if

$$H(Tx, Ty) \leq L \|x - y\|$$

for any  $x, y \in K$ ,  $L \geq 0$ .  $T$  is *nonexpansive* if  $L = 1$  and a *contraction* if  $L < 1$ .

$T$  is said to be *demiclosed* (on the convex subset  $K$ ) if  $x_n$  converges weakly to  $x$ ,  $y_n$  converges to  $y$ ,  $y_n \in Tx_n$ , imply  $y \in Tx$ .

$T$  is *semiconvex* (on  $K$ ) if for any  $x, y \in K$ ,  $z = \lambda x + (1 - \lambda)y$  ( $0 \leq \lambda \leq 1$ ), and any  $u \in Tx$ ,  $v \in Ty$  there exists  $\eta \in Tz$  such that  $\|\eta\| \leq \max\{\|u\|, \|v\|\}$ .

We introduce now the geometric sets which play a fundamental role in our results. We set (for any  $x, y \in X$ )

$$\tau(x, y) := \lim_{t \rightarrow 0^+} \frac{\|x + ty\| - \|x\|}{t}.$$

The following properties of  $\tau$  are well known [8]

$$\tau(ax, y) = \tau(x, y) \quad (a > 0) \quad (1)$$

$$\tau(x, y) \leq \|y\| \quad (2)$$

$$\tau(x, bx + cy) = b\|x\| + c\tau(x, y) \quad (b \text{ real}, c \geq 0) \quad (3)$$

$$\tau(x, y + z) \leq \tau(x, y) + \tau(x, z) \quad (4)$$

$$\tau(x, y) = \langle x, y \rangle \|x\|^{-1} \quad (\text{in Hilbert spaces}).$$

Following [2] we define, for  $x, y \in X$ ,  $\varepsilon > 0$ ,

$$LS(x, y; K) := \{z \in K : \tau(x - z, y - x) < 0\}$$

$$LS(x, y, \varepsilon; K) := \{z \in K : \tau(x - z, y - x) < \varepsilon\}.$$

In [2] it is shown that, put

$$G(x, y; K) := \{z \in K : \|z - y\| \leq \|z - x\|\},$$

$X$  is strictly convex if and only if  $G(x, y; X) \subseteq LS(x, y; X)$  for any  $x, y \in X$ . This remains true if  $y$  is replaced by a subset  $A$  of  $X$ , i.e.,  $X$  is strictly convex if and only if  $G(x, A; X) \subseteq LS(x, A; X)$  for any  $x \in X$  and  $A \subseteq X$ , where  $G$  and  $LS$  are so defined:

$$LS(x, A; K) := \{z \in K : \exists a \in A : \tau(x - z, a - x) < 0\} = \bigcup_{a \in A} LS(x, a; K)$$

$$LS(x, A, \varepsilon; K) := \{z \in K : \exists a \in A : \tau(x - z, a - x) < \varepsilon\} = \bigcup_{a \in A} LS(x, a, \varepsilon; K)$$

$$G(x, A; K) := \{z \in K : \exists a \in A : \|z - a\| \leq \|z - x\|\} = \bigcup_{a \in A} G(x, a; K).$$

## 2. NONEXPANSIVE MAPPINGS

Our first result extends [2, Theorem 1.3] concerning self-mappings to multivalued mappings.

**THEOREM 1.** *Let  $X$  be a real Banach space whose bounded closed convex subsets have the fixed point property for multivalued nonexpansive point-compact self-mappings.*

*Let  $K$  be a closed convex subset of  $X$  and  $T: K \rightarrow \mathcal{K}(K)$  nonexpansive. If there exists  $x_0 \in K$  such that  $LS(x_0, \overline{\text{co}}(Tx_0); K)$  is bounded, then  $T$  has a fixed point in  $K$ .*

*Prof.* Assume  $x_0 \notin Tx_0$  (otherwise we are done). It is enough to show that a bounded closed convex subset of  $K$  exists which is invariant under  $T$ .

We set

$$R := 4 \sup \{ H(\{z\}, \overline{\text{co}}(Tx_0)) : z \in LS(x_0, \overline{\text{co}}(Tx_0); K) \}$$

$$S := \{ z \in K : \exists v \in \overline{\text{co}}(Tx_0) \text{ such that } \|z - v\| \leq R \}.$$

Then one can verify that  $S$  is a nonempty bounded closed convex subset of  $K$ . It remains to show that  $T(S) \subseteq S$ . We consider separately the two cases

$$z \in S \cap LS(x_0, \overline{\text{co}}(Tx_0); K) \quad \text{and} \quad z \in S, z \notin LS(x_0, \overline{\text{co}}(Tx_0); K).$$

Assume first that  $z \in S \cap LS(x_0, \overline{\text{co}}(Tx_0); K)$  and let  $\eta \in \overline{\text{co}}(Tx_0)$ ,  $\eta \neq x_0$ . Then

$$\begin{aligned} \tau\left(x_0 - \frac{x_0 + \eta}{2}, \eta - x_0\right) &= \tau\left(\frac{x_0 - \eta}{2}, \eta - x_0\right) = (\text{from (1)}) \\ &= \tau(x_0 - \eta, \eta - x_0) = (\text{from (3)}) = -\|x_0 - \eta\| < 0, \end{aligned}$$

i.e.,  $(x_0 + \eta)/2 \in LS(x_0, \overline{\text{co}}(Tx_0); K)$ , and by definition of  $R$

$$\frac{R}{4} \geq H\left(\left\{\frac{x_0 + \eta}{2}\right\}, \overline{\text{co}}(Tx_0)\right) \geq \left\|\frac{x_0 + \eta}{2} - \eta\right\| = \frac{1}{2}\|x_0 - \eta\|,$$

i.e.,  $\|x_0 - \eta\| \leq R/2$ . Moreover,  $\|z - \eta\| \leq R/4$  since  $z \in LS(x_0, \overline{\text{co}}(Tx_0); K)$ , so

$$\|z - x_0\| \leq \|z - \eta\| + \|\eta - x_0\| \leq \frac{R}{4} + \frac{R}{2} < R.$$

Now [9, Lemma 2] assures that  $H(\overline{\text{co}}(Tz), \overline{\text{co}}(Tx_0)) \leq H(Tz, Tx_0)$  and so, from nonexpansivity of  $T$ , we have

$$H(\overline{\text{co}}(Tz), \overline{\text{co}}(Tx_0)) \leq \|z - x_0\| < R.$$

By definition of Hausdorff metric  $H$  it follows that for any  $u \in Tz$  there exists  $v \in \overline{\text{co}}(Tx_0)$  such that

$$\|u - v\| \leq H(\overline{\text{co}}(Tz), \overline{\text{co}}(Tx_0)) < R,$$

i.e.,  $Tz \subseteq S$ .

Assume now that  $z \in S$  but  $z \notin LS(x_0, \overline{CO}(Tx_0); K)$ . Then for any  $\eta \in \overline{CO}(Tx_0)$  we have

$$\tau(x_0 - z, \eta - x_0) \geq 0$$

and, since  $z \in S$ , there exists  $v \in \overline{CO}(Tx_0)$  such that  $\|z - v\| \leq R$ . Moreover,

$$\begin{aligned} \|z - x_0\| &= \tau(x_0 - z, x_0 - z) \quad (\text{from (3)}) \\ &= \tau(x_0 - z, v - z) - \tau(x_0 - z, v - x_0) \quad (\text{from (3)}) \\ &\leq \tau(x_0 - z, v - z) \\ &\leq \|v - z\| \quad (\text{from (2)}) \\ &\leq R. \end{aligned}$$

Finally from

$$H(\overline{CO}(Tz), \overline{CO}(Tx_0)) \leq H(Tz, Tx_0) \leq \|z - x_0\| \leq R$$

follows, as before, that for any  $u \in Tz$  there exists  $v \in \overline{CO}(Tx_0)$  such that

$$\|u - v\| \leq H(\overline{CO}(Tz), \overline{CO}(Tx_0)) \leq R,$$

i.e.,  $Tz \subseteq S$ , concluding the proof. ■

The next theorem extends [3, Theorem 3] to multivalued case.

**THEOREM 2.** *Let  $K$  be a closed convex subset of a real Banach space  $X$ , and let  $T: K \rightarrow \mathcal{K}(X)$  be a nonexpansive mapping which satisfies the following “inwardness” condition:  $T(x) \subseteq \overline{I_K(x)}$  for any  $x \in K$ , where  $I_K(x) := \{x + c(u - x) : u \in K, c \geq 1\}$  is the “inward set” of  $x$  relative to  $K$ . Suppose for some bounded set  $W \subseteq K$  that the set*

$$LS(W, TW; K) := \bigcap_{w \in W} LS(w, Tw, K)$$

*is bounded. Then there exists a bounded sequence  $\{x_n\} \subseteq K$  such that  $d(x_n, Tx_n) \rightarrow 0$  as  $n \rightarrow \infty$ .*

*Proof.* Let  $y \in K$  fixed, and for any  $\alpha \in (0, 1)$  define  $T_\alpha: K \rightarrow \mathcal{K}(X)$  by

$$T_\alpha(x) := (1 - \alpha)y + \alpha Tx.$$

It is easy to verify that  $T_\alpha$  is a contraction that satisfies the “inwardness condition”  $T_\alpha(x) \subseteq \overline{I_K(x)}$  for any  $x \in K$  (indeed, by convexity of  $K$  convexity of  $I_K(x)$  follows for any  $x \in K$ ).

Thus from [10, Theorem 3.4]  $T_\alpha$  has a fixed point  $x_\alpha \in K$ .

Suppose, by contradiction, that the set  $\{x_\alpha : \alpha \in (0, 1)\}$  is unbounded. Then it is possible to choose  $k \in (0, 1)$  such that

$$\sup_{w \in W} H(\{y\}, Tw) < d(x_k, W) \quad (5)$$

$$\sup_{z \in LS(W, TW; K)} \|z\| < \|x_k\|. \quad (6)$$

We will prove that  $x_k \in LS(W, TW; K)$ , i.e., that for any  $w \in W$  there exists  $\eta_w \in Tw$  such that  $\tau(w - x_k, \eta_w - w) < 0$ . Indeed, by  $x_k \in (1 - k)y + kTx_k$ , it follows that  $x_k = (1 - k)y + k\xi$ ,  $\xi \in Tx_k$ .

From nonexpansivity of  $T$  there exists  $\eta_w \in T(w)$  such that

$$\|\xi - \eta_w\| \leq \|x_k - w\| \quad (7)$$

and so

$$\begin{aligned} \tau(w - x_k, \eta_w - w) &= \tau(w - x_k, \eta_w - w + x_k - x_k) \\ &= -\|w - x_k\| + \tau(w - x_k, \eta_w - x_k) \quad (\text{from (3)}) \\ &\leq -\|w - x_k\| + \|\eta_w - x_k\| \quad (\text{from (2)}) \\ &= -\|w - x_k\| + \|\eta_w - (1 - k)y - k\xi \pm k\eta_w\| \\ &\leq -\|w - x_k\| + k\|\xi - \eta_w\| + (1 - k)\|\eta_w - y\| \\ &\leq (k - 1)\|x_k - w\| + (1 - k)\|\eta_w - y\| \quad (\text{from (7)}) \\ &\leq (k - 1)\|x_k - w\| + (1 - k)H(\{y\}, Tw) \\ &< (k - 1)\|x_k - w\| + (1 - k)d(x_k, W) \quad (\text{from (5)}) \\ &\leq (k - 1)\|x_k - w\| + (1 - k)\|x_k - w\| = 0, \end{aligned}$$

i.e.,  $x_k \in LS(W, TW; K)$  contradicting (6).

Thus  $M := \sup\{\|x_\alpha - y\|, \alpha \in (0, 1)\} < \infty$  and moreover

$$\begin{aligned} d(x_\alpha, Tx_\alpha) &= \inf_{z \in Tx_\alpha} \|x_\alpha - z\| \leq \left\| x_\alpha - \frac{x_\alpha - (1 - \alpha)y}{\alpha} \right\| \\ &= \frac{1 - \alpha}{\alpha} \|x_\alpha - y\| \leq \frac{1 - \alpha}{\alpha} M \end{aligned}$$

and this last converges to 0 as  $\alpha \rightarrow 1$ . ■

The following corollary improves [11, Corollary 2].

**COROLLARY 3.** *Let  $X, K, T$  be as in Theorem 2. Suppose that, for some bounded set  $W \subseteq K$  and  $\varepsilon > 0$ , the set*

$$LS(W, TW, \varepsilon; K) := \bigcap_{w \in W} LS(w, Tw, \varepsilon; K)$$

*is relatively compact. Then  $T$  has a fixed point.*

*Proof.* Let  $y \in K$  fixed and  $A := \sup\{\|y - \xi\|, \xi \in TW\}$ . Then for any  $\alpha$  belonging to

$$I := \left( \frac{A - \varepsilon}{A}, 1 \right) \cap (0, 1)$$

the mapping defined by  $T_\alpha(x) := (1 - \alpha)y + \alpha Tx$  is a contraction such that  $T_\alpha(x) \subseteq \overline{I_K(x)}$  and so, as in Theorem 2,  $T_\alpha$  has a fixed point  $x_\alpha \in T_\alpha(x_\alpha)$ .

We show that  $x_\alpha \in LS(W, TW, \varepsilon; K)$  for any  $\alpha \in I$ .

Indeed, since  $x_\alpha \in T_\alpha(x_\alpha)$ , for any  $w \in W$  there exists  $\eta_w \in Tw$  such that

$$\|x_\alpha - \eta_w\| \leq H(T_\alpha(x_\alpha), Tw). \quad (8)$$

Hence,

$$\begin{aligned} \tau(w - x_\alpha, \eta_w - w) &= \tau(w - x_\alpha, \eta_w - x_\alpha + x_\alpha - w) \\ &= -\|x_\alpha - w\| + \tau(w - x_\alpha, \eta_w - x_\alpha) \quad (\text{from (3)}) \\ &\leq -\|x_\alpha - w\| + \|\eta_w - x_\alpha\| \quad (\text{from (2)}) \\ &\leq -\|x_\alpha - w\| + H(T_\alpha(x_\alpha), Tw) \quad (\text{from (8)}) \\ &\leq -\|x_\alpha - w\| + H(T_\alpha(x_\alpha), T_\alpha(w)) + H(T_\alpha(w), Tw) \\ &\leq -\|x_\alpha - w\| + \alpha \|x_\alpha - w\| + H(T_\alpha(w), Tw) \\ &\quad (\text{from contractivity of } T_\alpha) \\ &= (\alpha - 1) \|x_\alpha - w\| + H(T_\alpha(w), Tw) \\ &\leq H(T_\alpha(w), Tw). \end{aligned}$$

Now, it is a routine calculation to verify that  $H(T_\alpha(w), Tw) \leq (1 - \alpha)A$ , and so, for the choice of  $\alpha$ , we obtain  $\tau(w - x_\alpha, \eta_w - w) < \varepsilon$ , i.e.,  $x_\alpha \in LS(W, TW, \varepsilon; K)$ .

By Theorem 2 we have  $d(x_\alpha, Tx_\alpha) \rightarrow 0$  for  $\alpha \rightarrow 1$ .

Let now  $\{x_{\alpha(n)}\}$  be a sequence of  $\{x_\alpha\}$  such that  $x_{\alpha(n)} \rightarrow x_0$  (such a sequence necessarily exists from relative compactness of  $LS(W, TW, \varepsilon; K)$ ). Then

$$\frac{x_{\alpha(n)}}{\alpha(n)} \rightarrow x_0, \quad \frac{x_{\alpha(n)} - (1 - \alpha(n))y}{\alpha(n)} \in Tx_{\alpha(n)}, \quad \frac{x_{\alpha(n)} - (1 - \alpha(n))y}{\alpha(n)} \rightarrow x_0$$

and from upper semicontinuity of  $T$  follows  $x_0 \in Tx_0$ . ■

**COROLLARY 4.** Let  $X$  be a reflexive real Banach space,  $K \subseteq X$  closed convex,  $T: K \rightarrow \mathcal{K}(X)$  a nonexpansive mapping such that  $T(x) \subseteq \overline{I_K(x)}$  for any  $x \in K$ ,  $I - T$  demiclosed on  $K$ .

Suppose for some bounded set  $W \subseteq K$  that the set  $LS(W, TW; K)$  is bounded. Then  $T$  has a fixed point.

*Proof.* As in the proof of Theorem 2 there exists a bounded sequence  $\{x_n\}$  such that

$$x_n = (1 - \alpha_n)y + \alpha_n y_n$$

with  $y_n \in Tx_n$ ,  $y_n$  bounded also, and  $\|x_n - y_n\| = (1 - \alpha_n)\|y_n\| \rightarrow 0$  as  $n \rightarrow \infty$ .

From reflexivity of  $X$  there exists a subsequence  $x_{n(j)}$  of  $\{x_n\}$  such that  $x_{n(j)}$  converges weakly to  $z$ , for some  $z \in K$ . So, since  $x_n - y_n \in (I - T)(x_n)$  and  $x_n - y_n \rightarrow 0$ , the demiclosure of  $I - T$  assures that  $0 \in (I - T)(z)$ , i.e.,  $z \in Tz$ . ■

The next corollary improves [11, Theorem 1].

**COROLLARY 5.** Let  $X, K, T, W$  be as in Theorem 2 and suppose that there exists  $\varepsilon > 0$  such that  $LS(W, TW, \varepsilon; K)$  is relatively weakly compact.

If  $I - T$  is demiclosed or semiconvex on  $K$ , then  $T$  has a fixed point.

*Proof.* First Case.  $I - T$  demiclosed.

As the proof of Corollary 4, since in this case  $\{x_n\} \subseteq LS(W, TW, \varepsilon; K)$  (see Corollary 3).

Second Case.  $I - T$  semiconvex.

We know (Corollary 3) that there exists a sequence  $\{x_n\} \subseteq LS(W, TW, \varepsilon; K)$  such that  $d(x_n, Tx_n) \rightarrow 0$ .

From [11, Propositions 1 and 2] it follows that for any  $r > 0$  the sets

$$H_r := \{x \in K : d(x, Tx) \leq r\}$$

are (nonempty) closed convex. Hence  $H_r$  are weakly closed for  $r > 0$ . Moreover the sets

$$\tilde{H}_r := H_r \cap \overline{LS(W, TW, \varepsilon; K)}^{\text{weak}}$$

are nonempty and weakly closed in  $\overline{LS(W, TW, \varepsilon; K)}^{\text{weak}}$  and obviously have the finite intersection property.

Therefore, by the weak compactness of  $\overline{LS(W, TW, \varepsilon; K)}^{\text{weak}}$  we have  $\bigcap \tilde{H}_r \neq \emptyset$ . It is clear that any point in  $\bigcap \tilde{H}_r$  is a fixed point of  $T$ . ■

**COROLLARY 6.** Let  $X, K, T, W, LS(W, TW; K)$  be as in Theorem 2. If  $X$  is reflexive and satisfies Opial's condition (i.e., if  $z_n$  converges weakly to  $z$  and  $z \neq v$  then  $\liminf_{n \rightarrow \infty} \|z_n - z\| < \liminf_{n \rightarrow \infty} \|z_n - v\|$ ), then  $T$  has a fixed point.



*Proof.* Immediate from Corollary 4 since if  $X$  satisfies Opial's condition and  $T$  is nonexpansive, then  $I - T$  is demiclosed on  $K$  by a result of [12]. ■

**COROLLARY 7.** *Let  $X, K, T, W, LS(W, TW; K)$  be as in Theorem 2. If  $X$  is uniformly convex, then  $T$  has a fixed point.*

*Proof.* Immediate from Corollary 4 since if  $X$  is uniformly convex and  $T$  is nonexpansive, then  $I - T$  is demiclosed on  $K$  by a result of [13]. ■

### 3. LIPSCHITZIAN PSEUDO-CONTRACTIVE MAPPINGS

In this last section we give a result concerning lipschitzian pseudocontractive mappings, where we say that  $T: K \rightarrow \mathcal{CB}(X)$  is pseudocontractive if for any  $x, y \in K, u \in Tx, v \in Ty, r > 0$  we have

$$\|x - y\| \leq \|(1 + r)(x - y) - r(u - v)\|.$$

This class of mappings includes, in the singlevalued case, all nonexpansive mappings, but for multivalued case this inclusion is not verified.

For example, the multivalued map  $T: \mathbb{R} \rightarrow \mathcal{K}(\mathbb{R})$  defined by  $T(x) = [x, x + 1]$  is nonexpansive but not pseudocontractive. (Here  $\mathbb{R}$  denotes the real numbers.)

The results concerning the existence of fixed points for pseudocontractive mappings are closely related with the existence of zeros of accretive mappings (see Morales [14]) since  $T$  is pseudocontractive if and only if  $I - T$  is accretive. The study of accretive mappings is very important in connection with the existence theory for nonlinear equations of evolution in Banach spaces (see, for example, [15–18]).

Theorem 1 (concerning nonexpansive multivalued self-mappings) is used in the proof of the following (concerning lipschitzian pseudocontractive multivalued mappings).

**THEOREM 8.** *Let  $X$  be a real Banach space whose bounded closed convex subsets have the fixed point property for multivalued nonexpansive point-compact self-mappings.*

*Let  $K$  be a closed convex subset of  $X$  and let  $T: K \rightarrow \mathcal{K}(X)$  be a lipschitzian pseudocontractive mapping which satisfies the inwardness condition  $Tx \subseteq \overline{I_K}(x)$  for any  $x \in K$ .*

*Suppose that there exist  $x_0 \in K$  and  $\varepsilon > 0$  such that  $LS((x_0, \overline{\text{co}}(Tx_0), \varepsilon; K)$  is bounded.*

*If  $TK \cap B_r$  is compact for any  $B_r := \{z \in K: \|z\| \leq r\}$  then  $T$  has a fixed point.*

*Proof.* Let  $L$  be the Lipschitz constant of  $T$  and select  $\alpha$  such that

$$0 < \alpha < \min(L^{-1}, \varepsilon(2LH(\{x_0\}, Tx_0) + L\varepsilon)^{-1}).$$

Then for any  $y \in K$  the mapping  $T_y: K \rightarrow \mathcal{K}(X)$  given by  $T_y(x) := (1 - \alpha)y + \alpha Tx$  is a contraction that satisfies the inwardness condition  $T_y(x) \subseteq \overline{I_K(x)}$  for any  $x \in K$ .

Hence, by [10, Theorem 3.4] the set  $F_\alpha(y)$  of fixed points of  $T_y$  is non-empty, closed and it is easy to verify that

$$F_\alpha(y) \subseteq (1 - \alpha)y + \alpha TF_\alpha(y) \quad (9)$$

$$y \in F_\alpha(y) \quad \text{iff} \quad y \in Ty. \quad (10)$$

Moreover we will show that for  $u, v \in K$  fixed

$$\|a - b\| \leq \|u - v\| \quad \text{for any } a \in F_\alpha(u), b \in F_\alpha(v). \quad (11)$$

Indeed  $a \in F_\alpha(u)$  and  $b \in F_\alpha(v)$  imply

$$a = (1 - \alpha)u + \alpha\eta, \quad \eta \in Ta \quad (12)$$

$$b = (1 - \alpha)v + \alpha\xi, \quad \xi \in Tb \quad (13)$$

and so, from pseudo-contractivity of  $T$ , from (12) and (13) and choosing  $r < \alpha/(1 - \alpha)$ ,

$$\begin{aligned} \|a - b\| &\leq \|(1 + r)(a - b) - r(\eta - \xi)\| \\ &= \left\| (1 + r)(a - b) - \frac{r}{\alpha} (a - (1 - \alpha)u - b + (1 - \alpha)v) \right\| \\ &= \left\| (1 + r)(a - b) - \frac{r}{\alpha} ((a - b) - (1 - \alpha)(u - v)) \right\| \\ &= \left\| \left(1 + r - \frac{r}{\alpha}\right) (a - b) - \frac{r}{\alpha} (1 - \alpha)(u - v) \right\| \\ &\leq \left(1 + r - \frac{r}{\alpha}\right) \|a - b\| + \frac{r}{\alpha} (1 - \alpha) \|u - v\|, \end{aligned}$$

i.e.,  $r(1 - \alpha)\alpha^{-1} \|a - b\| \leq r(1 - \alpha)\alpha^{-1} \|u - v\|$  proving (11).

It follows, in particular, that  $F_\alpha(y)$  belongs to  $\mathcal{CB}(K)$  for any  $y \in K$ . Besides  $TF_\alpha(y)$  is bounded (since  $T$  is lipschitzian) and by (9) we can conclude, under the hypothesis  $TK \cap B_r$  is compact for any  $r$ , that for any  $y \in K$ ,

$$F_\alpha(y) \in \mathcal{K}(K). \quad (14)$$

Now we note that

$$H(F_\alpha(x), F_\alpha(y)) \leq \sup_{a \in F_\alpha(x), b \in F_\alpha(y)} \|a - b\|$$

and so from (11) and (14) it follows that the mapping

$$F_\alpha: K \rightarrow \mathcal{K}(K), \quad x \rightarrow F_\alpha(x)$$

is nonexpansive and by (10) has the same fixed points of  $T$ .

Hence the claim will be proved if we are able to show that  $F_\alpha$  has a fixed point.

From Theorem 1 it is enough to prove that the set  $LS(x_0, \overline{\text{co}}(F_\alpha(x_0)); K)$  is bounded. For this we will see that

$$LS(x_0, \overline{\text{co}}(F_\alpha(x_0)); K) \subseteq LS(x_0, \overline{\text{co}}(Tx_0), \varepsilon; K). \quad (15)$$

Indeed, from (9) it follows that

$$\overline{\text{co}}(F_\alpha(x_0)) \subseteq (1 - \alpha)x_0 + \alpha \overline{\text{co}}(TF_\alpha(x_0))$$

and so  $z \in LS(x_0, \overline{\text{co}}(F_\alpha(x_0)); K)$  implies

$$\begin{aligned} z &\in LS(x_0, (1 - \alpha)x_0 + \alpha \overline{\text{co}}(TF_\alpha(x_0)); K) \\ &= \bigcup_{y \in \overline{\text{co}}(TF_\alpha(x_0))} LS(x_0, (1 - \alpha)x_0 + \alpha y; K) \\ &= \bigcup_{y \in \overline{\text{co}}(TF_\alpha(x_0))} \{z \in K : \tau(x_0 - z, (1 - \alpha)x_0 + \alpha y - x_0) < 0\} \\ &= \bigcup_{y \in \overline{\text{co}}(TF_\alpha(x_0))} \{z \in K : \tau(x_0 - z, \alpha(y - x_0)) < 0\} \\ &= \bigcup_{y \in \overline{\text{co}}(TF_\alpha(x_0))} \{z \in K : \tau(x_0 - z, y - x_0) < 0\} \quad (\text{from (3)}) \\ &= \bigcup_{y \in \overline{\text{co}}(TF_\alpha(x_0))} LS(x_0, y; K) \\ &= LS(x_0, \overline{\text{co}}(TF_\alpha(x_0)); K). \end{aligned}$$

Hence to prove (15) it is enough to prove that

$$LS(x_0, \overline{\text{co}}(TF_\alpha(x_0)); K) \subseteq LS(x_0, \overline{\text{co}}(Tx_0), \varepsilon; K). \quad (16)$$

First we note that

$$H(\overline{\text{co}}(TF_\alpha(x_0)), \overline{\text{co}}(Tx_0)) < \frac{\varepsilon}{2}. \quad (17)$$

Indeed

$$\begin{aligned}
 & H(TF_x(x_0), Tx_0) \\
 & \leq LH(F_x(x_0), \{x_0\}) \quad (\text{from [9, Theorem 2]}) \\
 & = L \sup_{a \in F_x(x_0)} \|x_0 - a\| \\
 & \leq L\alpha \sup_{a \in TF_x(x_0)} \|a - x_0\| \quad (\text{since } F_x(x_0) - x_0 \subseteq \alpha(TF_x(x_0)) - x_0) \\
 & = L\alpha H(TF_x(x_0), \{x_0\}) \\
 & \leq L\alpha(H(TF_x(x_0), Tx_0) + H(Tx_0, \{x_0\}))
 \end{aligned}$$

so that

$$H(TF_x(x_0), Tx_0) \leq \frac{L\alpha}{1 - L\alpha} H(Tx_0, \{x_0\}) < \frac{\varepsilon}{2} \quad (\text{by the choice of } \alpha).$$

From this (17) follows since

$$H(\overline{\text{co}}(TF_x(x_0)), \overline{\text{co}}(Tx_0)) \leq H(TF_x(x_0), Tx_0)$$

(see [9, Lemma 2]).

Finally we prove (16). Let  $z \in LS(x_0, \overline{\text{co}}(TF_x(x_0)); K)$ . Then there exists  $a \in \overline{\text{co}}(TF_x(x_0))$  such that  $\tau(x_0 - z, a - x_0) < 0$ . From (17) it follows that there exists  $b \in \overline{\text{co}}(Tx_0)$  such that  $\|a - b\| < \varepsilon/2$  and so

$$\begin{aligned}
 \tau(x_0 - z, b - x_0) &= \tau(x_0 - z, b - a + a - x_0) \\
 &\leq \tau(x_0 - z, a - x_0) + \tau(x_0 - z, b - a) \quad (\text{from (4)}) \\
 &< \tau(x_0 - z, b - a) \leq \|b - a\| \quad (\text{from (2)}) \\
 &< \frac{\varepsilon}{2} < \varepsilon,
 \end{aligned}$$

i.e.,  $z \in LS(x_0, \overline{\text{co}}(Tx_0), \varepsilon; K)$ . ■

*Remark 9.* The previous theorem works also under the weaker hypothesis that  $T$  maps bounded sets into relatively compact sets instead of  $TK \cap B_r$  compact for any  $r > 0$ .

## REFERENCES

1. W. A. KIRK AND W. O. RAY, Fixed point theorem for mappings defined on unbounded sets in Banach spaces, *Studia Math.* **64** (1979), 127–138.
2. A. CARBONE AND G. MARINO, Fixed points and almost fixed points of nonexpansive maps in Banach spaces, *Riv. Mat. Univ. Parma (4)* **13** (1987), 385–393.

3. G. MARINO AND P. PIETRAMALA, Fixed points and almost fixed points for mappings defined on unbounded sets in Banach spaces, *Atti Sem. Mat. Fis. Univ. Modena*, to appear.
4. T. C. LIM, A fixed point theorem for multivalued nonexpansive mappings in a uniformly convex Banach space, *Bull. Amer. Math. Soc.* **80** (1974), 1123–1125.
5. K. GOEBEL AND T. KUCZUMOV, A contribution to the theory of nonexpansive mappings, *Bull. Calcutta Math. Soc.* **70** (1978), 355–357.
6. W. O. RAY, Zeros of accretive operators defined on unbounded sets, *Houston J. Math.* **5** (1979), 133–139.
7. T. E. WILLIAMSON, A geometric approach to fixed points of non-self mappings  $T: D \rightarrow X$ , *Contemp. Math.* **18** (1983), 247–253.
8. R. C. JAMES, Orthogonality and linear functionals in normed linear spaces, *Trans. Amer. Math. Soc.* **61** (1947), 265–292.
9. S. B. NADLER, JR., Multi-valued contractive mappings, *Pacific J. Math.* **30** (1969), 475–488.
10. S. REICH, Approximate selections, best approximations, fixed points and invariants sets, *J. Math. Anal. Appl.* **62** (1978), 104–113.
11. K. JANAGI, On some fixed point theorems for multivalued mappings, *Pacific J. Math.* **87** (1980), 233–240.
12. E. LAMI-DOZO, Multivalued nonexpansive mappings and Opial's condition, *Proc. Amer. Math. Soc.* **38** (1973), 286–292.
13. F. E. BROWDER, Semiccontractive and semiaccretive nonlinear mappings in Banach spaces, *Bull. Amer. Math. Soc.* **74** (1968), 660–665.
14. C. MORALES, Zeros for strongly accretive set-valued mappings, *Comment. Math. Univ. Carolin.* **27** (1986), 455–469.
15. F. E. BROWDER, Nonlinear operators and nonlinear equations of evolutions in Banach spaces, in "Proceedings Sympos. Pure Math.," Vol. 18, Part 2, Amer. Math. Soc., Providence, RI, 1976.
16. K. DEIMLING, Zeros of accretive operators, *Manuscripta Math.* **13** (1974), 365–374.
17. T. KATO, Nonlinear semigroups and evolution equation, *J. Math. Soc. Japan* **19** (1967), 508–520.
18. R. H. MARTIN, Differential equations on a closed subset of a Banach space, *Trans. Amer. Math. Soc.* **179** (1973), 399–414.